




# Beef cattle thermoregulation in response to naturally occurring heat stress on pasture

**Raluca Mateescu, H. Hamblen, A.M. Zolini, S. Dikmen, P.A. Oltenacu, M.A. Elzo, and P.J. Hansen**

**Department of  
Animal Sciences**

**UF** | UNIVERSITY of  
**FLORIDA**

# Climatic stress and beef cattle

- 
- Major limiting factor of production efficiency
    - In beef cattle in tropical and subtropical environments.
    - In dairy cattle throughout most of the world.
  - > **50%** cattle in the world – maintained in hot and humid environments
    - including ~ 40% of beef cows in US.
  - Substantial differences in thermal tolerance
    - Among breeds
    - Among animals within breeds
  - Indication of opportunities for **selective improvement.**

# In response to heat stress cattle will:

- Regulate internal **heat production**
  - Modulating basal metabolic rate
  - Changing: feed intake, growth, lactation, activity
- Regulate **heat exchange**
  - Increasing blood flow to the skin
  - Increasing evaporative heat loss through sweating & panting



# Research Populations

- UF Multibreed Angus x Brahman Herd
  - Summer 2015, 2017
  - **286 cows**: from 100% Brahman to 100% Angus

Breed Group	Angus %	Brahman %
1	100	0
2	75	25
3	62.5	37.5
4	50	50
5	25	75
6	0	100

- Brangus heifers, Seminole Tribe of Florida
  - Summer 2016, 2017
  - **1,500** two-year old heifers


# Internal Body Temperature

- Vaginal **temperature** at 5-min intervals for 5 days
- Air temperature and relative humidity on pasture – every 15 min.
- The temperature-humidity index (**THI**) was used to quantify heat stress and it was calculated as in Dikmen et al., 2008:


$$\text{THI} = (1.8 \times T_{\text{db}} + 32) - [(0.55 - 0.0055 \times \text{RH}) \times (1.8 \times T_{\text{db}} - 26)]$$

DS1922L iButton Temperature Logger -  
Maxim Integrated Products, 120 San  
Gabriel Drive, Sunnyvale, CA  
Range: -40°C to +85°C  
Resolution: 0.0625°C (11 bit) or 0.5°C (8 bit)

iButton



# Thermotolerance measurements

- 
- Vaginal **temperature** 15 min over 5 days
  - Environmental data: temperature, humidity, **THI**
  - **Sweating** rate
  - **Coat**: color, coat score, hair length & diameter
  - **Temperament**: chute and exit score
  - Body **condition** score
  - **Skin** biopsies: for histology & gene expression
  - **Weight gain** over the summer/fall
  - Rump fat and rib fat ultrasound
  - Subsequent **pregnancy** status
  - **250K** genotypes


# Body temperature variables



- **Low** THI: **74** and **76**
- **High** THI: **84 – 86**
- **Average** THI: **79 – 81**
- Vaginal temperature for each cow Low, High, Avg. = **average temp of all the 5-min** measurements when the cow was exposed to that respective THI.
- Diff THI: High-Low THI


# Effect of breed composition on body temp

Body Temperature (°C) under **High** THI (heat stress)



# Effect of breed composition on body temp

Body Temperature (°C) under **High** and **Low** THI



Trait	$h^2$
Temp Diff Hi-Low	0.27
Temp High	0.11
Temp Low	0.25

# Thermotolerance in Brangus

- 1,500 Brangus 2-year old heifers (2016-2017)



Score 1  
Excessively Smooth  
N = 526




Score 2  
Fairly Smooth  
N = 197

## Coat score


1. excessively smooth
2. fairly smooth
3. long coat
4. woolly
5. excessively woolly coat

Effect	Estimate	Std. err	t value	Pr >  t
Coat 1 vs 2	-0.097	0.021	-4.64	<.0001
Chute 1 vs 2	-0.047	0.019	-2.42	<b>0.015</b>
Exit 1 vs 2	0.011	0.020	0.57	0.567

# Skin histology




6mm biopsy in formalin => histology




## • Histological skin characteristics:

- Distance from the sweat gland top to the skin surface
- Distance from the sweat gland bottom to the skin surface
- Skin, epidermis, dermis thickness
- Sweat and sebaceous gland number
- Sweat and sebaceous gland area


# Effect of breed composition on sweat glands




## Sweat Gland area



# Distance from the top of the SG to the skin surface



# Conclusions



- **Climatic stress** - major limiting factor of production efficiency in beef cattle in tropical and subtropical environments.
  - Expected to increase due to climate change.
- Differences in **thermal tolerance** exist:
  - Opportunities for selective improvement.
- **Genomic tools** are needed to select replacement heifers or bulls with increased thermotolerance.
- Development of the “**cow of the future**” with high productivity and resistant to heat stress will be realized through use of **genomic selection**.

# Acknowledgments

## University of Florida

- Dr. Pete Hansen
- Dr. Mauricio Elzo
- Dr. Dwain Johnson
- Dr. Tracy Scheffler
- Dr. Jason Schaffler
- Dr. Serdal Dikmen
- Danny Driver
- Michelle Driver
- Joel Leal, Heather Hamblen, Sarah Flowers, Kaitlyn Sarlo, Mesfin Gobena, Zaira Estrada
- Eduardo Rodriguez
- Adriana Zolini, William Ortiz, Samantha Eifert, Lauren Peacock, Alexa Chiroussot

## Seminole Tribe of Florida

- Alex Johns
- Phillip Clark
- Sheri Holmes
- Bobby Yates



United States Department of Agriculture  
National Institute of Food and Agriculture



FLORIDA  
CATTLEMEN'S  
ASSOCIATION

## Financial Support

- USDA-NIFA Grant 2017-67007-26143
- UF Agricultural Experim. Station
- UF ANS Hatch Project
- Seminole Tribe of Florida
- Brangus Breeders Association
- Florida Beef Council
- Florida Cattlemen's Association





# Comments / Questions

